

FAT DIGESTING ENZYMES

LIPOPAN is a high potency combination of pancreatic lipolytic enzymes. Pancreatic lipase is the most important enzyme for the digestion of dietary fats referred to as hydrolysis. The more effectively fats are hydrolyzed in the intestinal tract, the better is the absorption of fats. Complete fat absorption in turn stabilizes the blood sugar levels, supplies the endocrine glands with the raw materials to produce steroid hormones such as estrogens and cortisone, facilitates the absorption of Vitamins A, D, E, and K, and maintains the quality of healthy tissue, especially the skin. If the pancreatic lipase enzymes are deficient, poor fat digestion can adversely affect the entire body.

INGREDIENTS:

Each capsule contains:		% Daily Value
In a proprietary blend of:	265 mg	*
Pancreatic enzymes		

*Daily Value not established

Other Ingredients: Gelatin (capsule) and calcium citrate.

No yeast, wheat, soy, corn, dairy, salt, sugar, colors, chemical additives, or preservatives.

INDICATORS:

LIPOPAN was developed to provide the highest quality and most effective pancreatic enzymes for the complete digestion of fats. When taken after meals, LIPOPAN will ensure adequate dietary fat assimilation. Regular use will eventually enable a complete replenishment of the pancreas stores of lipase. LIPOPAN is recommended when various indicators which signal poor fat digestion are present.

1. Extended transit time for meals
2. Stools that float or have oil droplets evident
3. Unusual appetite even if a substantial meal has been eaten
4. A need to consume fatty foods such as chips, ice cream, fries, fried foods, bacon, and fatty meats
5. Dry, flaky skin, or skin that has oily patches amid areas of dryness. Lipomas, sebaceous cysts, psoriasis, dry and cracking heels, and fissured palms
6. Hypoglycemic symptoms between adequate meals
7. Difficulty gaining weight accompanied by a large appetite
8. Difficulty losing weight. Excess weight is characteristically soft and loose
9. Menstrual irregularities, impotence combined with apathy
10. Weak mucous membranes, frequent colds or allergies, hardening of the arteries, high blood pressure, calcium deficiency syndromes, and poor clotting factor

DIRECTIONS:

As a dietary supplement 1 to 2 LIPOPAN should be taken after meals, or as directed by a physician.

COMBINATIONS:

BilePlex — when there is sluggish digestion with unusually prolonged fullness after a meal.

ColAid — to enhance gall bladder activity and improve the emulsification of dietary fats — needed with constipation and poor transit time

Enzyme F/S — to improve the quality of small intestine lipase activity by promoting healthy flora

Lecithin — to assist in the thinning of bile and the emulsification of fats

LipaLiv — improves the liver's ability to process and distribute fats for use as bile

Formulated by Dr. Norman Easley.