

MICROBAC 2™

ANTI-YEAST FORMULA

MICROBAC 2 is the first chewable anti-yeast formula ever developed. It was made chewable because clinical research has demonstrated that the primary focus for elimination of Candida is in the mouth, throat, and stomach. Any attempt to permanently eliminate gastro-intestinal Candida infestations must begin in these areas. MICROBAC 2 contains a highly selective strain of lactic acid-producing bacteria which inhibits the growth of yeast and candidiasis. It also restricts the growth of salmonella, enterobacterium, and E. coli. MICROBAC 2's task is to control intestinal yeast infestation, facilitate normalization of the entire intestinal environment, and assist colonization of the intestine by compatible organisms.

INDICATORS:

The distinctive strains of bacteria in MICROBAC 2 are effective for the following conditions:

1. Candidiasis
2. Food poisoning
3. Pharyngeal yeast infection and earaches
4. Post-antibiotic therapy
5. Intestinal allergies due to yeast toxicity
6. Sweet cravings due to yeast activity
7. Vaginal infections caused by intestinal candidiasis
8. Sinusitis
9. Sore throats, flus, colds, earaches
10. Lymph node swelling with colds and flu
11. As prevention for infections

INGREDIENTS:

Each chewable tablet contains:		% Daily Value
In a proprietary blend of:	172 mg	*
Acidophilus extract and enzymes		

*Daily Value not established

Other Ingredients: Dextrose, maltodextrose, and flavoring.

No corn, soy, salt, yeast, egg, whey, milk, colors, chemical additives, or preservatives.

Potency Guaranteed: All the bacteria in this product have been included at their most effective clinical potency, and will maintain this potency at room temperature for 1 year from purchase. Refrigeration will assist in maintaining this potency.

DIRECTIONS:

As a dietary supplement, begin by dissolving in mouth 1 MICROBAC 2 tablet 4 times a day, for 1-2 weeks on an empty stomach up to 15 minutes before eating, or at least 1 hour after meals and before bed, or as directed by a physician. Then decrease MICROBAC 2 to 1 tablet every AM and PM. For children 2 to 8 years of age the recommended dosage is half the adult dosage. In extreme cases, MICROBAC 2 may be required for as much as 8 weeks.

NOTE: For Candida see CANDIDA PROTOCOL

COMBINATIONS:

MicroBac 3 — to re-colonize the intestine allowing for sufficient bacterial replacement to assist in the control of "die-off" toxins

ReBuild — if a "leaky" intestinal lining exists enabling "die-off" to readily be absorbed
L-G-K-S or NuLiv — to assist the liver in processing "die-off" toxins (to prevent nausea, fatigue, skin breakouts, and the onset of illness)

Enzyme Pro or F/S — to facilitate more efficient digestion and to support bacterial colonization

Formulated by Dr. Norman Easley.